

What is the True Description of Charged Black Holes?

Gary T. Horowitz*

Department of Physics
University of California
Santa Barbara, CA 93106-9530
gary@cosmic.physics.ucsb.edu

Abstract

If string theory describes nature, then charged black holes are not described by the Reissner-Nordström solution. This solution must be modified to include a massive dilaton. In the limit of vanishing dilaton mass, the new solution can be found by a generalization of the Harrison transformation for the Einstein-Maxwell equations. These two solution generating transformations and the resulting black holes are compared. It is shown that the extremal black hole with massless dilaton can be viewed as the “square root” of the extremal Reissner-Nordström solution. When the dilaton mass is included, extremal black holes are repulsive, and it is energetically favorable for them to bifurcate into smaller holes.

* To appear in the Brill Festschrift, B. L. Hu and T. A. Jacobson, eds., Cambridge University Press, 1993.

1. Introduction

It is a pleasure to honor Dieter Brill on the occasion of his sixtieth birthday. Over the years, Dieter has worked on many aspects of general relativity. But two of his recent interests are negative energy (in higher dimensional theories) [1], and the possibility that extremal charged black holes can quantum mechanically bifurcate [2]. I would like to describe some recent work which touches on both of these areas.

For many years, it has been widely believed that static charged black holes in nature are accurately described by the Reissner-Nordström solution. This is the result of two powerful theorems: the uniqueness theorem [3] which proves that the only static black hole solution to the Einstein-Maxwell equations is Reissner-Nordström, and the no-hair theorem [4] which shows that if one adds simple additional matter fields to the theory, the only static black hole is still Reissner-Nordström. The extra fields either fall into the hole or radiate out to infinity. Although this solution seems physically reasonable in most respects, its extremal limit has two puzzling features. The first is that the event horizon is infinitely far away along a spacelike geodesic, but only a finite distance away along a timelike or null geodesic. The second is that the extremal black hole has zero Hawking temperature, but nonzero entropy as measured by its surface area.

Recently, a slightly different picture of charged black holes has emerged. This is the result of studying gravitational consequences of string theory. A new class of charged black hole solutions have been found¹ which do not have the puzzling features of Reissner-Nordström. In the extremal limit, the area of the event horizon of these new black holes goes to zero, which is consistent with zero entropy. From another standpoint (for a magnetically charged black hole) the horizon moves off to infinity in all directions: timelike, spacelike, and null. These solutions should be of particular interest to Dieter Brill since we will see that they are even more likely to bifurcate than the analogous black holes in the Einstein-Maxwell theory. In addition, although the total energy remains positive for these four dimensional holes, we will see that some of them resemble negative energy objects in that they are repulsive.

In addition to the metric $g_{\mu\nu}$ and Maxwell field $F_{\mu\nu}$, string theory predicts the existence of a scalar field called the dilaton. The dilaton has an exponential coupling to F^2 which implies that it cannot remain zero when $F^2 \neq 0$. This is the key difference from the previous no-hair theorems and shows that the charged black holes will differ from the

¹ For recent reviews, see [5][6].

Reissner-Nordström solution. Although the dilaton is massless while supersymmetry is unbroken, it is expected to become massive at low energies. The breaking of supersymmetry in string theory is a nonperturbative effect which is not well understood at this time. So we cannot yet calculate reliably the dilaton potential from first principles. However the qualitative behavior of the solutions can be obtained by considering the simplest possibility $m^2\phi^2$. We are thus led to consider the low energy action:

$$S = \int d^4x \sqrt{-g} [R - 2(\nabla\phi)^2 - 2m^2\phi^2 - e^{-2\phi}F^2] \quad (1.1)$$

with equations of motion

$$\nabla_\mu(e^{-2\phi}F^{\mu\nu}) = 0, \quad (1.2a)$$

$$\nabla^2\phi - m^2\phi = -\frac{1}{2}e^{-2\phi}F^2, \quad (1.2b)$$

$$G_{\mu\nu} = 2\nabla_\mu\phi\nabla_\nu\phi + 2e^{-2\phi}F_{\mu\rho}F_\nu{}^\rho - g_{\mu\nu}[(\nabla\phi)^2 + m^2\phi^2 + \frac{1}{2}e^{-2\phi}F^2] \quad (1.2c)$$

When $F_{\mu\nu} = 0$, these equations reduce to those of the Einstein-massive scalar field theory. The no-hair theorems show that the only black hole solutions are Schwarzschild with $\phi = 0$. However, as we just remarked, when the charge is nonzero, the dilaton will not be constant, and will alter the geometry. It does not appear possible to write the exact solution to (1.2) describing a static, spherically symmetric, charged black hole in closed form. But we will see that by combining general arguments, approximation methods, and numerical results, we can obtain a fairly complete understanding of their properties.

2. Solutions with a Massless Dilaton

Let us begin by considering the limit where the dilaton mass is negligible. It will be shown in the next section that this is appropriate for small black holes. Even with $m = 0$, the exponential coupling of the dilaton to the Maxwell field appears to prevent a simple closed form solution. However exact solutions can be found [7][8] which are, in some respects, even simpler than the Reissner-Nordström solution. In the Einstein-Maxwell theory there is a transformation, found by Harrison [9], which maps stationary vacuum solutions into stationary charged solutions. It turns out that there is a similar transformation for the theory (1.1) with $m = 0$ [10]. In this section, we compare these

two transformations and the resulting charged black holes. For simplicity, we will consider only static solutions.

We start with the familiar Einstein-Maxwell theory. Suppose

$$ds^2 = -\lambda dt^2 + h_{ij} dx^i dx^j \quad (2.1)$$

is a static vacuum solution. The Harrison transformation begins by rewriting this metric as

$$ds^2 = -\lambda dt^2 + \lambda^{-1} \gamma_{ij} dx^i dx^j \quad (2.2)$$

where $\gamma_{ij} = \lambda h_{ij}$. The charged solution is obtained by keeping γ_{ij} unchanged and setting

$$\tilde{\lambda} = \frac{\lambda}{(\cosh^2 \alpha - \lambda \sinh^2 \alpha)^2} \quad (2.3a)$$

$$A_t = \frac{(\lambda - 1) \sinh 2\alpha}{2(\cosh^2 \alpha - \lambda \sinh^2 \alpha)} \quad (2.3b)$$

where α is a free parameter. Clearly, $\alpha = 0$ leaves the solution unchanged. Since the factor in the denominator appears frequently, it is convenient to introduce the notation

$$e^{-2\phi} \equiv \cosh^2 \alpha - \lambda \sinh^2 \alpha \quad (2.4)$$

Notice that ϕ , at this point, has nothing to do with the dilaton. We are considering solutions to the Einstein-Maxwell equations and ϕ is simply defined by (2.4). Equation (2.3a) can now be rewritten $\tilde{\lambda} = \lambda e^{4\phi}$ and the new metric can be expressed

$$ds^2 = -\lambda e^{4\phi} dt^2 + e^{-4\phi} h_{ij} dx^i dx^j \quad (2.5)$$

Let us apply this transformation to the Schwarzschild solution:

$$\lambda = 1 - \frac{2M_0}{\rho}, \quad h_{ij} dx^i dx^j = \left(1 - \frac{2M_0}{\rho}\right)^{-1} d\rho^2 + \rho^2 d\Omega, \quad (2.6)$$

In this case, $e^{-2\phi} = 1 + 2M_0 \sinh^2 \alpha / \rho$. So the new metric becomes

$$ds^2 = - \left(1 - \frac{2M_0}{\rho}\right) \left(1 + \frac{2M_0 \sinh^2 \alpha}{\rho}\right)^{-2} dt^2 + \left(1 + \frac{2M_0 \sinh^2 \alpha}{\rho}\right)^2 \left[\left(1 - \frac{2M_0}{\rho}\right)^{-1} d\rho^2 + \rho^2 d\Omega \right] \quad (2.7)$$

This can be simplified by introducing a new radial coordinate $r = \rho + 2M_0 \sinh^2 \alpha$. Then

$$\rho(\rho - 2M_0) = (r - 2M_0 \sinh^2 \alpha)(r - 2M_0 \cosh^2 \alpha) \equiv (r - r_-)(r - r_+) \quad (2.8)$$

where we have defined $r_+ \equiv 2M_0 \cosh^2 \alpha$ and $r_- \equiv 2M_0 \sinh^2 \alpha$. The metric (2.7) now becomes:

$$ds^2 = -\frac{(r - r_+)(r - r_-)}{r^2} dt^2 + \frac{r^2}{(r - r_+)(r - r_-)} dr^2 + r^2 d\Omega \quad (2.9)$$

This is the familiar form of the Reissner-Nordström solution with the event horizon at $r = r_+$ and the inner horizon at $r = r_-$. Notice that the original Schwarzschild singularity $\rho = 0$ has become the inner horizon, while the Reissner-Nordström singularity $r = 0$ corresponds to a negative value of ρ in the original Schwarzschild metric. The Maxwell field is

$$F_{rt} = \frac{M_0 \sinh 2\alpha}{r^2} \quad (2.10)$$

which shows that the total charge is $Q = M_0 \sinh 2\alpha$. The total mass is $M = (r_+ + r_-)/2 = M_0 \cosh 2\alpha$. The extremal limit is obtained by taking $\alpha \rightarrow \infty, M_0 \rightarrow 0$ keeping $M_0 e^{2\alpha}$ finite. In this limit, the two horizons coincide $r_+ = r_-$. The event horizon becomes degenerate, with nonzero area but vanishing Hawking temperature.

Now we turn to the theory with a massless dilaton. There is a solution generating transformation directly analogous to the Harrison transformation. It is usually discussed in terms of the conformally rescaled metric $\hat{g}_{\mu\nu} = e^{2\phi} g_{\mu\nu}$ (where ϕ is now the dilaton) which is called the string metric. This is the metric that a string directly couples to. To avoid confusion, the metric we have been using until now will be called the Einstein metric (since it has the standard Einstein action). If one starts with a static vacuum solution with (string) metric of the form $\hat{ds}^2 = -\lambda dt^2 + h_{ij} dx^i dx^j$ and $\phi = 0$, then a new solution is obtained by keeping h_{ij} unchanged, and setting

$$\tilde{\lambda} = \frac{\lambda}{(\cosh^2 \alpha - \lambda \sinh^2 \alpha)^2} \quad (2.11a)$$

$$A_t = \frac{(\lambda - 1) \sinh 2\alpha}{2\sqrt{2}(\cosh^2 \alpha - \lambda \sinh^2 \alpha)} \quad (2.11b)$$

$$e^{-2\phi} = \cosh^2 \alpha - \lambda \sinh^2 \alpha \quad (2.11c)$$

The similarity to (2.3) is remarkable. The change in λ is identical to the Harrison transformation, and the expression for the vector potential A_t differs only by a factor of $\sqrt{2}$. In addition, the solution for the dilaton is exactly what we called ϕ in (2.4). (This is why we

chose that definition for ϕ .) To compare with the Reissner-Nordström solution, we should rescale back to the Einstein metric:

$$ds^2 = -\tilde{\lambda}e^{-2\phi}dt^2 + e^{-2\phi}h_{ij}dx^i dx^j \quad (2.12)$$

In terms of the original λ we get

$$ds^2 = -\lambda e^{2\phi}dt^2 + e^{-2\phi}h_{ij}dx^i dx^j \quad (2.13)$$

If one now compares (2.13) with the solution to the Einstein-Maxwell theory (2.5), one sees that *the only difference in the solutions with and without a massless dilaton is a factor of two in the exponent.*

Applying this transformation to the Schwarzschild solution (2.6), we obtain the metric

$$ds^2 = -\left(1 - \frac{2M_0}{\rho}\right) \left(1 + \frac{2M_0 \sinh^2 \alpha}{\rho}\right)^{-1} dt^2 + \left(1 + \frac{2M_0 \sinh^2 \alpha}{\rho}\right) \left[\left(1 - \frac{2M_0}{\rho}\right)^{-1} d\rho^2 + \rho^2 d\Omega\right] \quad (2.14)$$

together with the Maxwell field and dilaton

$$A_t = -\frac{M_0 \sinh 2\alpha}{\sqrt{2}(\rho + 2M_0 \sinh^2 \alpha)} \quad (2.15a)$$

$$e^{-2\phi} = 1 + \frac{2M_0 \sinh^2 \alpha}{\rho} \quad (2.15b)$$

The factor of two difference in the exponent has the following important consequence: The surface $\rho = 0$ in the metric (2.14) still has zero area and does not become an inner horizon. The total mass is now $M = M_0 \cosh^2 \alpha$ and the charge is $Q = M_0 \sinh 2\alpha / \sqrt{2}$. Introducing the same radial coordinate as before, $r = \rho + 2M_0 \sinh^2 \alpha$, the solution takes the remarkably simple form

$$ds^2 = -\left(1 - \frac{2M}{r}\right) dt^2 + \left(1 - \frac{2M}{r}\right)^{-1} dr^2 + r \left(r - \frac{Q^2}{M}\right) d\Omega$$

$$F_{rt} = \frac{Q}{r^2} \quad e^{2\phi} = 1 - \frac{Q^2}{Mr} \quad (2.16)$$

Note that the metric in the $r - t$ plane is identical to Schwarzschild! (Although the mass is different from the one we started with.) There is an event horizon at $r = 2M$ and no

inner horizon. The only difference from Schwarzschild is that the area of the spheres is reduced by an amount depending on the charge. This area goes to zero when $r = Q^2/M$ ($\rho = 0$) resulting in a curvature singularity.

Clearly, the solution (2.16) describes a black hole only when the singularity is inside the event horizon i.e. $Q^2 < 2M^2$. Linearized perturbations about this solution have been studied [11] and remain well behaved outside the horizon: the black hole is stable. In the extremal limit, $Q^2 = 2M^2$, the event horizon shrinks down to zero area and becomes singular. The resulting spacetime describes neither a black hole (with a spacelike singularity), nor a conventional naked singularity (which is timelike). Since the causal structure in the $r - t$ plane is independent of Q , the extremal black hole has a null singularity. Its Penrose diagram is identical to the region $r > 2M$ of Schwarzschild, with both the future and past horizons, $r = 2M$, replaced by singularities.

The fact that the area of the event horizon goes to zero in the extremal limit has two important consequences. First, it follows immediately from the area theorem that there is no classical process by which a nearly extremal black hole can become extremal [12]. This is also true for the Reissner-Nordström solution, but the proof is more involved. Second, if one interprets the area as a measure of the entropy of a black hole through the Hawking-Bekenstein formula, $S = A/4$, then the extremal black holes have zero entropy. This is what one might expect for a ground state. The nonzero area of the extremal Reissner-Nordström solution has always been puzzling from this standpoint, since it seems to indicate a highly degenerate ground state. Although this black hole resolves one puzzle, it creates a new one. For a static, spherically symmetric black hole, the Hawking temperature can be found by analytically continuing to imaginary time and computing what periodicity is necessary to avoid a conical singularity. Since this only involves the $r - t$ part of the metric which is identical to Schwarzschild for the solution (2.16), the Hawking temperature is also identical: $T = 1/8\pi M$, independent of Q ! In particular, the Hawking temperature does not go to zero as one approaches the extremal limit. This leads one to worry that the black hole might continue to evaporate past the extremal limit to form a true naked singularity. However, the backreaction clearly becomes important as one approaches the extremal limit, and a complete calculation is necessary to understand the evolution².

In terms of the original Schwarzschild mass M_0 and transformation parameter α , the extremal limit again corresponds to $M_0 \rightarrow 0, \alpha \rightarrow \infty$ keeping $M_0 e^{2\alpha}$ fixed. In this limit,

² For a review of recent work in this area, see [13].

$\lambda \rightarrow 1$ and $h_{ij} \rightarrow \delta_{ij}$. So the extremal black hole metrics (2.5) and (2.13) are determined entirely by the function $e^{2\phi}$. Since the solutions with and without a dilaton differ only by a factor of two in the exponent, *one can view the extremal black hole with dilaton as the square root of the extremal Reissner-Nordström solution*. It is not clear what the physical interpretation of this is. On the one hand, the statement is clearly coordinate dependent. It holds in isotropic coordinates where the spatial metric is manifestly conformally flat. On the other hand, the statement applies not just to single black holes but extends to multi-black hole solutions as well. In the Einstein-Maxwell theory, since there is no force between extremal black holes (with the same sign of the charge), there exist static, multi-black hole solutions. These are the Majumdar-Papepetrou solutions [14] and take the form

$$ds^2 = -e^{4\phi} dt^2 + e^{-4\phi} d\vec{x} \cdot d\vec{x} \quad (2.17)$$

with

$$e^{-2\phi} = 1 + \sum_i \frac{2M_i}{|\vec{x} - \vec{x}_i|} \quad (2.18)$$

In exact analogy, the extremal black holes with a massless dilaton also have no force between them. The fact that $Q^2 = 2M^2$ rather than $Q^2 = M^2$ in the extremal limit means that there is a stronger electrostatic repulsion which exactly cancels the additional attractive force due to the dilaton. The multi-black hole solution turns out to be simply the “square root” of (2.17):

$$ds^2 = -e^{2\phi} dt^2 + e^{-2\phi} d\vec{x} \cdot d\vec{x} \quad (2.19)$$

with $e^{-2\phi}$ again given by (2.18).

The solution generating transformations (2.3) and (2.11) produce solutions with electric charge. But one can easily obtain solutions with magnetic charge by applying a duality transformation. When the dilaton is present, the duality transformation corresponds to replacing $F_{\mu\nu}$ and ϕ with $\tilde{F}_{\mu\nu} \equiv \frac{1}{2}e^{-2\phi}\epsilon_{\mu\nu}{}^{\lambda\rho}F_{\lambda\rho}$ and $\tilde{\phi} = -\phi$. This leaves the stress energy tensor invariant and hence does not change the Einstein metric. But since the dilaton changes sign, it does change the string metric. For electrically charged black holes, the string metric is obtained by multiplying (2.19) by $e^{2\phi}$. The result is that the spatial metric is completely flat and g_{tt} is identical to the Majumdar-Papepetrou solutions. On the other hand, since ϕ changes sign under duality, the string metric describing several magnetically charged black holes is obtained by multiplying (2.19) by $e^{-2\phi}$ where ϕ is still given by (2.18). The resulting metric has $g_{tt} = -1$ and a spatial metric which is identical to the

Majumdar-Papapetrou solution. In other words, the string metric describing several extremal black holes is obtained from the analogous Einstein-Maxwell solution by flattening either the time or space part of the metric (depending on the type of charge) leaving the rest unchanged.

The magnetically charged case is of particular interest. Recall that in the spatial part of the Reissner-Nordström metric, the horizon moves off to infinity as one approaches the extremal limit. The extremal geometry resembles an infinite throat attached to an asymptotically flat region. Since $g_{tt} = -1$ for the string solution, the horizon is now infinitely far away in timelike and null directions as well as spacelike. It has been suggested [15] that this type of geometry may play an important role in explaining what happens to the information that falls into a black hole, after it evaporates.

3. Solutions with a Massive Dilaton

We now wish to include the effects of the dilaton mass.³ Unfortunately, exact black hole solutions do not seem to be expressible in closed form. Given the simple form of the Reissner-Nordström (2.9) and massless dilaton (2.16) solutions, we will assume a metric of the form

$$ds^2 = -\lambda dt^2 + \lambda^{-1} dr^2 + R^2 d\Omega \quad (3.1)$$

where λ and R are functions of r only.

We first consider the asymptotic form of the solution. We are interested in solutions that are asymptotically flat, which requires that $\phi \rightarrow 0$ at infinity. For large r , the right hand side of the dilaton equation (1.2b) behaves like Q^2/r^4 , and the derivative term becomes negligible. The dilaton thus falls off like $|\phi| \sim Q^2/m^2 r^4$. Recall that a massless scalar field falls off as $1/r$ while a massive field with localized sources falls off exponentially. Here we have a massive field with a source that falls off polynomially, which results in the unusual asymptotic form of ϕ . Note that the limit $m \rightarrow 0$ keeping r fixed is not well behaved. This is because $|\phi| \sim Q^2/m^2 r^4$ only in the asymptotic region where r is large compared to the Compton wavelength of the dilaton, i.e. $rm \gg 1$.

Now consider the metric equation. At large distances, the Maxwell contribution to the stress tensor will be $O(1/r^4)$ while all terms involving the dilaton will fall off much faster.

³ This section is based work done in collaboration with J. Horne [16]. For another discussion of black holes with a massive dilaton see [17].

Thus, the asymptotic form of the field equation is identical to the Einstein-Maxwell theory, and the solution is just Reissner-Nordström. One can calculate the first order correction to the Reissner-Nordström solution by treating the dilaton terms as a perturbation and one finds

$$\lambda \sim 1 - \frac{2M}{r} + \frac{Q^2}{r^2} - \frac{Q^4}{5m^2r^6}, \quad (3.2a)$$

$$R \sim r \left(1 - \frac{2Q^4}{7m^4r^8} \right). \quad (3.2b)$$

Although the value of the dilaton mass is not known, one can place a lower limit from the fact that the $1/r^2$ force law has been confirmed down to scales of about 1 cm. This requires that the dilaton Compton wavelength be less than 1 cm or $m > 10^{-5}\text{eV}$. It follows that for a solar mass black hole⁴, $Mm > 10^5$. In other words, the black hole is much larger than the Compton wavelength of the dilaton. In this case, the deviation from Reissner-Nordström remains small until one is well inside the event horizon: At the horizon, $\delta\lambda < (Q/M)^4(Mm)^{-2} \ll 1$ and $\delta R < (Q/M)^4(Mm)^{-4} \ll 1$. So outside the horizon, the solution will be very similar to Reissner-Nordström. Since $\phi \sim Q^2/m^2r^4 < (Q/M)^2(Mm)^{-2} \ll 1$, the dilaton remains small everywhere outside the event horizon. For $Q \approx M$, the inner horizon of Reissner-Nordström is close to the event horizon and the corrections due to the dilaton will still remain small there. So unlike the massless dilaton solution, a large black hole coupled to a massive dilaton can have two horizons. This is what one should expect physically. When $Mm \gg 1$, the dilaton is essentially stuck in the bottom of its potential well and does not affect the solution significantly.

Even though the dilaton does not qualitatively affect the geometry, it has an important consequence near the extremal limit. Like Reissner-Nordström, the horizons will coalesce in the extremal limit. However, the condition for when this occurs is no longer exactly $Q^2 = M^2$. The new condition is found by asking when λ has a double zero. If we substitute the unperturbed condition, $Q^2 = M^2$ and $r = M$, into the correction term in (3.2a) we find that to first order in the dilaton, $\lambda = 0$ is equivalent to

$$r^2 - 2Mr + \left(Q^2 - \frac{1}{5m^2} \right) = 0 \quad (3.3)$$

⁴ In geometrical units, the dilaton mass m has dimensions of inverse length and is related to the inverse Compton wavelength. On the other hand, the black hole mass M has dimensions of length and is related to the size of the black hole. Thus Mm is dimensionless. Alternatively, one can view all quantities as dimensionless and measured in Planck units.

This will have a double zero when

$$M^2 = Q^2 - \frac{1}{5m^2} \quad (3.4)$$

So $M^2 < Q^2$ in the extremal limit, just like the massless dilaton solutions. But at infinity, we have seen that the solution always approaches the Reissner-Nordström solution, for which objects with $Q > M$ are repulsive. Since Q^2 is strictly greater than M^2 in the extremal limit, nonextremal black holes can also have a charge greater than their mass. We conclude that *nearly extremal and extremal black holes with a massive dilaton are repulsive*. Roughly speaking, this is a result of the fact that the presence of the dilaton near the horizon allows $Q^2 > M^2$ but the dilaton mass cuts off the attractive dilaton force at large separations. This appears to be the first example of gravitationally bound repulsive objects.

Since the extremal black hole has a degenerate horizon, its Hawking temperature vanishes. However, this does not guarantee that it is stable. Extremal black holes might quantum mechanically bifurcate. In the context of the Einstein-Maxwell theory, this was considered by Brill [2]. He found an instanton describing the splitting of an extremal Reissner-Nordström throat, and argued that there should exist a similar finite action instanton describing the splitting of the extremal Reissner-Nordström black hole. This has also been considered in the theory with a massless dilaton [18]. However in both of these theories, the total mass M of the black hole is proportional to its charge Q in the extremal limit. Thus one black hole of charge Q has the same mass as n black holes of charge Q/n . The solutions are degenerate. This is not the case for the theory with a massive dilaton. It is now energetically favorable for an extremal black hole to bifurcate. From (3.4), the mass of a single extremal black hole of charge Q is $M_1 = (Q^2 - 1/5m^2)^{\frac{1}{2}}$ while the mass for n widely separated extremal black holes with charge Q/n is

$$M_n = n \left[\left(\frac{Q}{n} \right)^2 - \frac{1}{5m^2} \right]^{\frac{1}{2}} = \left[Q^2 - \frac{n^2}{5m^2} \right]^{\frac{1}{2}}. \quad (3.5)$$

Clearly, M_n is a decreasing function of n . These black holes should be even more likely to bifurcate than the ones considered previously.

The above expression for M_n only applies to large black holes, since it is based on (3.4). (More precisely, it is valid when $mM_n/n \gg 1$.) However the conclusion is likely to hold more generally. This is because it follows from the fact that extremal black holes

are repulsive, which in turn is a consequence of two general properties of the extremal solutions: they always approach Reissner-Nordström asymptotically and have $Q^2 > M^2$. Thus extremal black holes will probably continue to bifurcate until they have a single unit of charge. At this point the black hole is small compared to the Compton wavelength of the dilaton, $Mm \ll 1$. (Black holes of this type can also be obtained by starting with a large hole with small charge, and letting it evaporate. The extremal limit will not be reached until $Mm \ll 1$.) For these small black holes there is a region $M \ll r \ll 1/m$ which is far from the black hole but inside the dilaton Compton wavelength. One can solve the dilaton equation (1.2b) exactly for $r \gg M$ since the spacetime is essentially flat. The result is that for $M \ll r \ll 1/m$, ϕ behaves like a massless dilaton: $\phi \propto 1/r$. So one expects the black holes to behave qualitatively like the massless dilaton solution discussed in the previous section. Numerical calculations confirm that this is indeed the case [16]. In particular, there is no inner horizon and in the extremal limit, the event horizon shrinks to zero area and becomes singular. The string metric describing an extremal magnetically charged black hole has an infinite throat.

We have seen that black holes with a massive dilaton can be viewed as interpolating between the Reissner-Nordström solution (when their mass is large), and the massless dilaton solution (when their mass is small). One can say something about the transition region $Mm \approx 1$ by considering when a degenerate horizon can exist. If there is a radius r_0 for which λ and λ' both vanish, the field equations yield the following condition on the value of ϕ at r_0

$$\frac{e^{2\phi}}{\phi(1+\phi)^2} = Q^2 m^2. \quad (3.6)$$

The left hand side has a minimum of $e^2/4$ when $\phi = 1$. So for $Q^2 m^2 < e^2/4$, there can not exist a degenerate horizon. For $Q^2 m^2 > e^2/4$ there are two possible values of ϕ , but the equations of motion also imply that $\lambda'' \propto (1 - \phi)$ so one needs $\phi < 1$ to have $\lambda'' > 0$ as required for an event horizon. At the critical value, $Q^2 m^2 = e^2/4$, λ'' also vanishes yielding a triple horizon. This suggests that there may be solutions with three distinct horizons. But at the moment, there is no direct evidence for this.

These intermediate size black holes can have an unusual property when described in terms of the string metric. One can show that the radius of the spheres of spherical symmetry in the Einstein metric, R , is monotonically increasing outside the horizon as expected. In the string metric, the radius of the spheres is Re^ϕ . For an electrically charged hole, it turns out that ϕ is also increasing outside the horizon so the spheres indeed

become larger as r increases. However, for a magnetically charged hole, ϕ changes sign and is monotonically decreasing outside the horizon. So Re^ϕ need not be an increasing function of r . It certainly increases near infinity, and for the massless dilaton solution (2.16), it is monotonically increasing everywhere. However, for nonzero dilaton mass, one can show that it decreases outside the horizon for nearly extremal black holes with $Mm \approx 1$. This means that the spatial geometry contains a wormhole. Unlike the familiar wormhole in the maximally extended Schwarzschild solution, this wormhole is outside the horizon. It is static and transversable.

Combining this with our previous results, we see that the string metric describing an extremal, magnetically charged black hole has a rather exotic dependence on mass. It resembles the Reissner-Nordström metric with a degenerate horizon when $Mm \gg 1$. A wormhole forms outside the degenerate horizon when $Mm \approx 1$. Finally, the horizon and wormhole both disappear and are replaced by an infinite throat when $Mm \ll 1$.

4. Conclusions

We have discussed two related issues. On the mathematical side, we have seen that despite the apparent complication introduced by a massless dilaton, the theory shares some of the features of the Einstein-Maxwell theory. In particular, there is a solution generating transformation which is remarkably similar to the Harrison transformation. The resulting extremal black hole can be viewed as the “square root” of the extremal Reissner-Nordström solution. On the physical side, we have seen that the evolution of charged black holes predicted by string theory is quite different from standard lore. Black holes initially (when $Mm \gg 1$) are quite similar to the Reissner-Nordström solution. But as they evolve (through Hawking evaporation and possible quantum bifurcation) they reach a stage where $Mm \ll 1$. At this point, the black holes are similar to the massless dilaton solutions. For a dilaton mass of $m = 1\text{TeV}$, the transition will occur at a black hole mass of $M = 10^{11}\text{gms}$ which is well above the Planck scale. This ensures that Planck scale corrections, which have been neglected throughout, will still be negligible.

What is the most likely endpoint of this evolution? Assuming the charge is not completely radiated away, one expects to be left with an extremal black hole with unit charge⁵. Since charge is quantized, this must be stable against further bifurcation. These

⁵ Although likely, this is still uncertain due to the fact that the Hawking temperature does not vanish as one approaches the extremal limit of these small black holes.

remnants would be pointlike objects, with charge greater than their mass, which repel each other. In other words, they would appear very much like elementary particles. (See also [11].) In fact, by shifting the dilaton potential so that its minimum is at $\phi_0 \neq 0$, one can change the charge to mass ratio of the extremal black holes. There even exists a value of ϕ_0 such that the extremal black hole of unit charge has the mass of an electron. In the past, one of the main objections to interpreting an electron as a black hole was the fact that its charge is much larger than its mass (in geometrical units) so that according to the Reissner-Nordström solution, it would be a naked singularity. The massive dilaton predicted by string theory removes this objection and opens up the possibility of a much closer connection between elementary particles and black holes.

Acknowledgments

It is a pleasure to thank my collaborators D. Garfinkle, A. Strominger, and especially J. Horne. This work was supported in part by NSF Grant PHY-9008502.

References

- [1] D. Brill and H. Pfister, “States of Negative Total Energy in Kaluza-Klein Theory”, *Phys. Lett.* **B228**, 359 (1989); D. Brill and G. Horowitz, “Negative Energy in String Theory”, *Phys. Lett.* **B262**, 437 (1991).
- [2] D. Brill, “Splitting of an Extremal Reissner-Nordström Throat via Quantum Tunneling”, *Phys. Rev.* **D46**, 1560 (1992).
- [3] W. Israel, “Event Horizons in Static Electrovac Spacetimes”, *Commun. Math. Phys.* **8**, 245 (1968).
- [4] J. Chase, *Commun. Math. Phys.* **19**, 276 (1970); J. Bekenstein, *Phys. Rev.* **D5**, 1239 (1972).
- [5] G. Horowitz, “The Dark Side of String Theory: Black Holes and Black Strings”, to appear in the proceedings of the 1992 Trieste Spring School on String Theory and Quantum Gravity, hep-th/9210119.
- [6] A. Sen, “Black Holes and Solitons in String Theory”, Tata preprint TIFR-TH-92-57, hep-th/9210050.
- [7] G. Gibbons, “Antigravitating Black Hole Solitons with Scalar Hair in N=4 Supergravity”, *Nucl. Phys.* **B207**, 337 (1982); G. Gibbons and K. Maeda, “Black Holes and Membranes in Higher-Dimensional Theories with Dilaton Fields”, *Nucl. Phys.* **B298**, 741 (1988).
- [8] D. Garfinkle, G. Horowitz and A. Strominger, “Charged Black Holes in String Theory,” *Phys. Rev.* **D43**, 3140 (1991); **D45**, 3888(E) (1992).
- [9] B. Harrison, “New Solutions of the Einstein-Maxwell Equations from Old”, *J. Math. Phys.* **9**, 1744 (1968).
- [10] S. Hassan and A. Sen, “Twisting Classical Solutions in Heterotic String Theory,” *Nucl. Phys.* **B375**, 103 (1992); J. Maharana and J. Schwarz, “Noncompact Symmetries in String Theory, Caltech preprint CALT-68-1790, hep-th/9207016.
- [11] C. Holzhey and F. Wilczek, “Black Holes as Elementary Particles,” *Nucl. Phys.* **B380**, 447 (1992).
- [12] D. Garfinkle, private communication.
- [13] J. Harvey and A. Strominger, “Quantum Aspects of Black Holes”, to appear in the proceedings of the 1992 Trieste Spring School on String Theory and Quantum Gravity, hep-th/9209055.
- [14] S. Majumdar, *Phys. Rev.* **72**, 930 (1947); A. Papepetrou, *Proc. Roy. Irish Acad.* **A51**, 191 (1947); J. Hartle and S. Hawking, *Commun. Math. Phys.* **26**, 87 (1972).
- [15] T. Banks, A. Dabholkar, M. Douglas, and M. O’Loughlin, “Are Horned Particles the Climax of Hawking Evaporation?”, *Phys. Rev.* **D45**, 3607 (1992); T. Banks and M. O’Loughlin, “Classical and Quantum Production of Cornucopions at Energies Below 10^{18} GeV”, Rutgers preprint RU-92-14, hep-th/9206055; T. Banks, M. O’Loughlin,

- and A. Strominger, “Black Hole Remnants and the Information Puzzle”, Rutgers preprint RU-92-40, hep-th/9211030.
- [16] J. Horne and G. Horowitz, “Black Holes Coupled to a Massive Dilaton”, Nucl. Phys. B to appear, hep-th/9210012.
- [17] R. Gregory and J. Harvey, “Black Holes with a Massive Dilaton”, Enrico Fermi Preprint EFI-92-49, hep-th /9209070.
- [18] R. Kallosh, A. Linde, T. Ortin, A. Peet, and A. Van Proeyen, “Supersymmetry as a Cosmic Censor”, Stanford preprint SU-ITP-92-13; R. Kallosh, T. Ortin, and A. Peet, “Entropy and Action of Dilaton Black Holes”, Stanford preprint SU-ITP-92-29, hep-th/9211015.